



UPDATES ON IMMUNOTHERAPY FOR PRIMARY BRAIN TUMORS

LISA FELDMAN, MD, PHD

Assistant Professor, Department of Surgery
Division of Neurosurgery
City of Hope National Medical Center

Disclosures




- I have nothing to disclose.

Glioblastoma (GBM)




- GBMs are aggressive gliomas, notoriously invasive, rendering complete surgical resection impossible.
- Remains the most common brain tumor in adults, less than 10% patients live beyond 5 years from diagnosis, with median survival less than 20 months.
 - With surgery, chemotherapy, alternating electrical fields, radiation (Alexander & Cloughesy, 2017; Stupp et al., 2017).
- Even in children and young adults, brain tumors remain the leading cause of cancer-related deaths (Pollack, Agnihotri, & Broniscer, 2019)
- Because the blood-brain barrier (BBB) precludes the transmission of many standard therapies, Neuro-oncologists are desperate for novel therapies.

GBM




Dec 2002



March 2003

Estimated % of deaths and new cases 2020



Estimates are rounded to the nearest 10, and cases exclude basal cell and squamous cell skin cancers and in situ carcinoma except urinary bladder. Estimates do not include Puerto Rico or other US territories. Ranking is based on modeled projections and may differ from the most recent observed data.

©2020, American Cancer Society, Inc., Surveillance Research


According to the Central **Brain Tumor** Registry of the United States (CBTRUS), it is **estimated** that **84,170 new** cases of **primary malignant** and non-malignant brain and other CNS tumors will be **diagnosed** in the United States in 2021.

(CBTRUS, 2020)


Gliomas: W.H.O. Grading System




<u>Anaplastic Features</u>	<u>Grade</u>	<u>Incidence</u>
None	I	Rare
Nuclear atypia	II	~15%
Mitosis	III	~30%
Vascular endothelial proliferation	IV	50-60%
Necrosis	IV	



Grade III Astrocytoma



Anaplastic Oligodendroglioma




**Grade IV Astrocytoma
Glioblastoma (GBM)**

MGMT (O⁶-Methylguanine-DNA Methyl Transferase)



- MGMT is an enzyme that repairs damage to DNA from alkylating agents.
- Methylation (inactivation) of the MGMT gene results in lower levels of MGMT in tumor.
- **Methylated MGMT** occurs in 50% of GBM pts
 - A prognostic factor for GBM patients
 - A predictive factor for response to treatment with an alkylating agent (temozolomide, BCNU)



(Hegi et al., 2005)

Select Biomarkers in Glioblastoma




Biomarker	Prognostic Indication	Prognostic Association	
		Favorable	Poor
<i>MGMT</i> methylation^[1]	<ul style="list-style-type: none"> Methylated in 30%–60% of cases Methylated <i>MGMT</i> increases response to chemotherapy Unmethylated <i>MGMT</i> decreases response to chemotherapy 	✓	
<i>IDH1/2</i> mutations^[2,3]	<ul style="list-style-type: none"> More common in lower grade glial tumors <i>IDH1/2</i> mutation occurs in approximately 3.7% of primary GBMs versus 73.3% in secondary GBM 	✓	
<i>EGFR</i> amplification^[4,5]	Observed in ~50% of primary glioblastomas		✓
<i>EGFRvIII</i> mutation^[4]	<ul style="list-style-type: none"> <i>EGFR</i>-amplified cells often contain <i>EGFRvIII</i> mutation, which confers constitutive activity 30% glioblastoma tumors express <i>EGFRvIII</i> 		✓

(Johnson et al., 2012; McNamara, Sahebjam, & Mason, 2013; Nobusawa, Watanabe, Kleihues, & Ohgaki, 2009; Preusser et al., 2011; Stupp et al., 2014; Yan et al., 2009)



Standard Treatments for Glioblastoma



- **No curative options!**
 - **Goal: Preserve QOL, neuro function and extending survival** (Henriksson, Asklund, & Poulsen, 2011)
- **Surgery**
 - Pathological diagnosis
 - Maximal safe resection can increase survival
 - Alleviation of mass effect (pressure)
 - Reverse or prevent neurological deficit
 - Avoid causing neurological deficits



(Dandy, 1928)



Standard Treatments for Glioblastoma

- **Radiation**


- For 6 weeks, focal brain RT
- Side effects: hair loss, fatigue, cognitive decline

- **Chemotherapy**

- **Temozolomide (TMZ):** oral and IV alkylating agent (during RT and for at least 6 months afterwards)
- Alkylating agents bind to DNA, impair replication/transcription → cell death (Fu, Calvo, & Samson, 2012)
- TMZ crosses BBB and is spontaneously converted into active metabolite in CNS (Agarwala & Kirkwood, 2000)

- **Toxicity**

- Most common: constipation, nausea
- Dose-limiting: neutropenia (14%), thrombocytopenia (19%)




(Wesolowski, Rajdev, & Mukherji, 2010)

Treatment Challenges



1. Brain tumors are invasive
2. Standard therapies (surgery, radiation, and chemotherapy) may cause brain injury
3. The blood-brain barrier prevents penetration of most chemotherapies into the brain
4. Tumor heterogeneity
5. Immunosuppressive microenvironment





Immunotherapy




- Tumor cells have cleverly evolved to evade their destruction by the host immune system.
- Immunotherapy (AKA biological therapy, biotherapy or biological response modifier therapy) stimulates the immune system to recognize and target cancer cells and/or lessen treatment-related side effects.
- Current immunotherapies are available for breast, hematological, kidney, lung, melanoma, mesothelioma, ovarian, pancreatic, prostate cancers....

Timeline of oncolytic virus development from glioblastoma therapy




- **Oncolytic virotherapy** employs viruses with a preserved potential for an active viral life cycle.
- The chief paradigm is tumor-selective conditional viral replication, resulting in lytic tumor cell destruction and release of thousands of viral progeny.
- Newly released viruses go on to infect neighboring tumor cells, theoretically causing and maintaining a wave of virus attack throughout the tumor.



Immune checkpoint inhibitors (ICIs):

- Immune checkpoints are molecules on certain immune cells that need to be activated or inactivated to start an immune response.
- Some tumors utilize ICIs to evade immune attack
- **Programmed cell death protein-1 (PD-1)** dampens anti-tumor immune responses (Kamath & Kumthekar, 2018; Simonelli et al., 2018)



© 2015 Terese Winslow LLC
U.S. Govt. has certain rights

Chimeric antigen receptor (CAR) T cells



- CAR T cells were designed over four decades ago to overcome an ineffective immune system by genetically modifying T lymphocytes to recognize and eliminate cancer cells (Gross, Waks, & Eshhar, 1989).
- In this adoptive cell therapy, cells are harvested from patients, modified to target specific proteins expressed by the tumor, and re-introduced into the patient with the goal of destroying tumor cells.
- Shown great promise in blood-borne cancers such as relapsed or refractory B-cell lymphomas by targeting CD19 (Maude et al., 2018; Park et al., 2018).
- Currently FDA-approved for relapsed or refractory diffuse large B-cell lymphoma (DLBCL) and acute lymphoblastic lymphoma (ALL).
 - Detected in CSF

Limitations of CAR T therapy for Glioma



- Tumor heterogeneity:
 - CAR T cells are particularly successful at targeting and destroying B-cell malignancies because these T cells are engineered to bind to a single molecule that is uniformly expressed on the surface of all B-cell derived tumors, CD19.
 - GBM tumors are notorious for having [intra-tumor](#), and [inter-tumor heterogeneity](#) of cellular, genetic and molecular signatures (Brown, Styli, Kaye, & Mantamadiotis, 2019; Neftel et al., 2019; Verhaak et al., 2010).

GBM “Harsh” Microenvironment



- GBM tumors actively contribute to immune suppression through a host of well-orchestrated strategies to create a T-cell suppressive tumor microenvironment.
 - Microenvironment suppress their activity and proliferation by **expressing inhibitory cell-surface molecules** (such as PD-L1, CD95) (Ricklefs et al., 2018) or by **releasing immunosuppressive tumor-derived soluble factors and cytokines** (such as prostaglandin E2, IL6, IL10 and TGFβ) (Hao et al., 2002).
- Microenvironment preferentially promotes trafficking of suppressive cell populations, such as regulatory T cells (Tregs), tumor-associated macrophages, microglia and myeloid-derived suppressor cells (Choi, Maus, June, & Sampson, 2019; Dubinski et al., 2016; Hussain et al., 2006) and creates other physical and metabolic blockades (Mirzaei, Sarkar, & Yong, 2017; Newick, O'Brien, Moon, & Albelda, 2017).

What if the therapy works TOO well...




- **Cytokine release syndrome (CRS)**, or the systemic elevation in several cytokines including **IL-6 and IFN-γ**, is a common toxicity associated with CD19-CAR T cells that is indicative of immunotherapeutic potency (Lee et al., 2014).
- Neurotoxicity (encephalopathy, delirium, seizures, edema) is also a common treatment-related toxicity of CD19-CAR therapy, resulting from increased inflammatory cytokine levels and endothelial dysfunction of the blood-brain barrier (Gust and Turtle, 2017; Santomasso and Brenjens 2018; Gauthier and Turtle, 2018)
- **Avoiding any severe brain inflammation and severe CRS is of utmost importance in GBMs, as increased intracranial pressure in patients already with increased mass effect from tumor can lead to deadly outcomes.**
- CAR T cell trials in GBM have thus far reported **less severe** CRS and neurotoxicity-like adverse events as compared to CAR T cells targeting hematological cancers.
- To give Dex or not??

CRS Treatments



- One strategy is to desensitize CAR T cells to steroids is to genetically disrupt the glucocorticoid receptor (Menger et al., 2015), while others use anti-IL6 antibody (O'Rourke et al., 2017) or Bevacizumab (Migliorini et al., 2018) to reduce local inflammation.
- A balance is needed in treating CAR T patients with symptomatic brain edema while prioritizing for CAR T cell therapeutic activity, and at City of Hope we do this by limiting dexamethasone to 6 mg in a 24-hour period in our patients.

Clinical Studies of CAR-T Cells for Glioblastoma



City of Hope


IL13Rα2+

NCT00730613 (2002-2008)
NCT01082926 (2011-2013)

NCT02208362 (2015-ongoing)
41BBζ-CAR Tmem

HER2+

NCT03389230 (2018-ongoing)
41BBζ-CAR Tmem



Baylor

HER2+

NCT02442297;
NCT01109095
CD28ζ-CAR T cells

EGFRvIII+

Upenn/UCSF

NCT02209376
41BBζ-CAR T
cell

NCI

NCT01454596
CD28-41BBζ-
CAR T cell
Flu/Cy; IL-2


Duke

NCT02664363;
NCT03283631
CD28-41BBζ-
CAR T cell






Clinicaltrials.gov

Interleukin 13 Receptor $\alpha 2$ (IL-13R $\alpha 2$)



- IL-13R $\alpha 2$ not only blocks the normal apoptotic pathway, but also induces upregulation of STAT3, promotes invasion and metastasis via extracellular signal-regulated kinase/activator protein 1, and promotes tumor immune escape via upregulation of transforming growth factor β (Thaci et al., 2014)
- Expressed in up to 75% GBMs with high variability (Jarboe, Johnson, Choi, Lonser, & Park, 2007; Joshi, Plautz, & Puri, 2000; Liu et al., 2000; Saikali et al., 2007)

CAR-T Cells Therapy for Glioblastoma: Laboratory to Clinic





Primary patient-derived gliomas incubated with autologous T cells
(captured over 8-12 hrs)

Clinical Evaluation of First-Generation Autologous IL13-zetakine T cells Locally injected for the Treatment of GBM





Patient Safety and Tolerability					
Patient	T cell Doses	Max Tolerated Dose	Cumulative Cell Dose	Serious Adverse Events (SAE) (therapy related probable)	Survival post-recurrence (months)
UPN028	11 of 12 (dose 7)	10 ⁸	9.6x10 ⁸	none	10.3
UPN031	12 of 12	10 ⁸	10.6x10 ⁸	Transient neurologic worsening (grade 3)	8.6
UPN033	12 of 12	10 ⁸	10.6x10 ⁸	none	13.9



Regression of Recurrent Multifocal GBM Following Intraventricular Delivery of IL13R α 2-CAR T cells




10 intraventricular (ICV) infusions



(Brown et al., 2018)

IL13R α 2-targeted CAR T immunotherapy for leptomeningeal disease clinical trial



Serial imaging: MRI brain with contrast, FDG-PET




Collection of peripheral blood and CSF followed by intrathecal injection of 100M cells IL13R α 2 CAR T cells

14 Current GBM clinical trials at COH



- IL13R α 2 CAR T cells with or without Nivolumab and Ipilimumab
- CAR T cells with Chlorotoxin Tumor-Targeting Domain
- IL13R α 2 CAR T cells for leptomeningeal disease
- Memory-Enriched T Cells in Treating Patients with Recurrent Grade III, IV glioma
- Genetically Modified T cells in patients with Recurrent malignant glioma

Acknowledgements



Behnam Badie, MD



Stephen Forman, MD




Christine Brown, PhD



**Vivian Chiu, Renate Starr
Cindy Xin**



**Aniee Sarkissian, Alfonso Brito
Brenda Aguilar**



Lihong Weng, Darya Alizadeh



**Julie Ostberg,
Jamie Wagner, Anita Kurien**

TCTRL Lab Members

- Wen-Chung Chang
- Vanessa Jonsson, PhD
- Jeff Kuo
- Saul Priceman, PhD
- Kirsten Rood, PhD
- Michalina Silva
- Sandra Thomas, PhD
- Robyn Wong
- Xiuli Wang, MD, PhD
- All Forman Lab Members

TCTRL Manufacturing

- Araceli Naranjo
- Jessica Chang
- Vidusha Cyril
- Maria-Isabel Garcia
- Alina Oancea
- Laurelin Wolfenden
- Jinny Paul
- Winnie Wong
- Joseph Done
- Leslie Reyes

COH Clinical Team

- Suzette Blanchard
- Fatemeh Bidmeshki, RN-BC
- Elizabeth Budde, MD, PhD
- Massimo D'Apuzzo, MD, PhD
- Kim Margolin, MD
- Samer Khaled, MD
- Julie Kilpatrick, CRN
- Leslie Popplewell, MD
- Jana Portnow, MD
- Julie Ann Ressler, MD
- Jennifer Simpson, CRC
- Yasmine Shad & QS
- Joseph Gold & CBG
- Alpha Clinic Team

COH Research Collaborators

- Margarita Gutova, MD
- Russell Rockne, PhD
- Jack Shively, PhD
- Tim Synold, PhD; CISCL
- John Williams, PhD

Other Collaborators

- Michael Jensen, MD
Seattle Children's Research Institute

Funding

- CIRM Early Translational Award
- CIRM CLIN2-10248
- CIRM Alpha Clinic
- Gateway for Cancer Research
- IVY Foundation
- NCI/FDA R01-FDA005129
- Norris Foundation
- Cancer Center Support Grant (P30CA33572)
- Mustang Bio, Inc Research Support

Thank You:
Patients and their families

