

**Multidisciplinary Approaches to Cancer Symposium** 

# Minimally Invasive Management of Pelvic Metastasis

John A. deVries, MD, MS

**Assistant Clinical Professor** 

Chief, Orthopedic Surgery

City of Hope



#### Disclosures

Consultant for Stryker

This presentation and/or comments will be free of any bias toward or promotion of the above referenced companies or their product(s) and/or other business interests.

This presentation and/or comments will provide a balanced, non-promotional, and evidence-based approach to all diagnostic, therapeutic and/or research related content.

This presentation has been peer-reviewed and no conflicts were noted.

### Cultural Linguistic Competency (CLC) & Implicit Bias (IB)

#### **STATE LAW:**

The California legislature has passed <u>Assembly Bill (AB) 1195</u>, which states that as of July 1, 2006, all Category 1 CME activities that relate to patient care must include a cultural diversity/linguistics component. It has also passed <u>AB 241</u>, which states that as of January 1, 2022, all continuing education courses for a physician and surgeon **must** contain curriculum that includes specified instruction in the understanding of implicit bias in medical treatment.

The cultural and linguistic competency (CLC) and implicit bias (IB) definitions reiterate how patients' diverse backgrounds may impact their access to care.

#### **EXEMPTION:**

Business and Professions Code 2190.1 exempts activities which are dedicated solely to research or other issues that do not contain a direct patient care component.

This presentation is dedicated solely to research or other issues that do not contain a direct patient care component.

#### Metastatic Bone Disease

- 20 million people with cancer by 2030
- 3<sup>rd</sup> most common site of metastasis (after liver and lung)
- 300-600,000 people in a given year with MBD
- 5-6% breast cancer patients present with MBD, increasing yearly
- Nearly 1/5 total cost of oncology care is from MBD
- Patients with MBD more than double their yearly treatment cost
- A skeletal related event can significantly increase costs and burden (up to 20x)
- Prophylactic surgery has been shown to decrease overall costs, decrease LOS, and increase functionality, and life expectancy vs. surgery after fracture.

DiCaprio, et al.





Figure 1 Cancer prevalence in the United States through the decades. Source of data: National Cancer Institute.

Table 1 Incidence of primary tumors that commonly metastasize to bone


Cancer type	Estimated number of new cases in the US, 2020	Incidence of MBD in patients with advanced disease (%)
Breast	279,100	65–75
Prostate	191,930	65–75
Lung	228,820	30-40
Renal	73,750	40
Thyroid	52,890	20–25
Bladder	81,400	60
Melanoma	100,350	14–45

MBD, metastatic bone disease.

## The Problem



## Acetabular Protrusio



# Cup Cage Construct





# Harrington Pelvic Reconstruction







There's an easier way.






# Minimally Invasive Stabilization with or without Ablation for Metastatic Periacetabular Tumors

D. Ian English, MD, MA, William B. Lea, MD, David M. King, MD, Sean M. Tutton, MD, FSIR, and John C. Neilson, MD

Investigation performed at the Division of Orthopaedic Surgery, Froedtert Hospital, Wauwatosa, and the Division of Interventional Radiology, Medical College of Wisconsin, Milwaukee Wisconsin



	All Patients (N = 38)			Patients without Other Surgical Treatments (N = 29)		
	Preoperative	Postoperative	P Value	Preoperative	Postoperative	P Value
Total	7.8	19.6	<0.001	7.9	19.7	<0.001
Pain	0.9	3.7	< 0.001	0.8	3.5	< 0.001
Emotional acceptance	1.1	4.2	< 0.001	1.0	4.2	< 0.001
Use of supports	1.1	2.3	< 0.001	1.1	2.2	0.0014
Walking ability	1.8	3.1	< 0.001	1.9	3.2	< 0.001
Gait	1.5	3.2	< 0.001	1.6	3.2	< 0.001
Function	1.5	3.2	< 0.001	1.4	3.0	< 0.001

<sup>\*</sup>Musculoskeletal Tumor Society Scoring system; a score of 0 to 5 is assigned in each of 6 categories, which are then combined for a maximum possible score of 30.


Narcotic use‡		
Reduced	81% (26 of 32)	87% (20 of 23)
No change	16% (5 of 32)	9% (2 of 23)
Increased	3% (1 of 32)	4% (1 of 23)

#### Minimally Invasive Stabilization with or without Ablation for Metastatic Periacetabular Tumors


D. Ian English, MD, MA, William B. Lea, MD, David M. King, MD, Sean M. Tutton, MD, FSIR, and John C. Neilson, MD

Investigation performed at the Division of Orthopaedic Surgery, Froedtert Hospital, Wauwatosa, and the Division of Interventional Radiology, Medical
College of Wisconsin, Milwaukee Wisconsin


#### Preparation for AORIF: Partial insertion of Cannulated Screws



#### Ablation




#### Osteoplasty with a balloon



#### Reinforcement with bone cement



#### Internal Fixation




Minimally Invasive Image-Guided Ablation, Osteoplasty, Reinforcement, and Internal Fixation (AORIF) for Osteolytic Lesions in the Pelvis and Periarticular Regions of Weight-Bearing Bones 🦠 🔁

Francis Y. Lee MD, PhD, Igor Latich MD, Courtney Toombs MD, Alana Mungur MD, Devin Conway MD, Kareme Alder BS, Izuchukwu Ibe MD, Dieter Lindskog MD and Gary Friedlaender MD Journal of Vascular and Interventional Radiology, 2020-04-01, Volume 31, Issue 4, Pages 649-658.e1, Copyright © 2019 SIR


#### Pain Reduction and Function

- Lee et al. 26 patients, VAS scores 8.32 -> 2.36, statistical improvement of MSTS score
- Hirase et al. Systemic review 579 patients (mostly cement alone) 74.2% significant pain and function improvement
- Lozano-Calderon et al. 39 patients with photodynamic intrapelvic nails VAS 8 -> 0 at 6 weeks.

#### Mechanical Studies









Pelvis weight-bearing ability after minimally invasive stabilizations for periacetabular lesion

William B. Lea<sup>1</sup> | Sean M. Tutton<sup>1</sup> | Naif Alsaikhan<sup>1</sup> | John C. Neilson<sup>2</sup> | Sebastian Schafer<sup>3</sup> | David M. King<sup>2</sup> | Mei Wang<sup>2,4</sup> |

### Case #1









42-year-old female with metastatic breast cancer progressing after RT/chemo

# Intra Operative Ablation and Planning





# Post Operative Course








17

Pre-op walker and 8/10 pain -> 1/10, no pain meds, no limp, 23 hr hospital stay

### Minimally Invasive Photodynamic Nails - Illuminoss




### Case #2






73-year-old female, metastatic renal cell, wheelchair bound






### Case #3



51 yoF bedbound, pelvic breast cancer Mets, S/p RT









### Case #4












41-year-old female, metastatic breast, inability to bear weight







CITY OF HOPE

26

### Case #5



66-year-old female with metastatic lung and sacral insufficiency fracture, progressive low back pain



VAS pain 10 to 2 on POD1







Define Pa







#### Who is a candidate?

- Any patient with painful pelvic or sacral metastasis
  - o Myeloma, metastatic renal, thyroid, breast, lung, prostate
  - o Even stress or insufficiency fractures if minimally displaced
- Specifically pain bearing weight, usually periacetabular, usually lytic lesion
- In conjunction with Radiation Therapy studies ongoing
- Best patients have good preoperative function, and early diagnosis of metastasis
- Contra indicated in displaced fracture open surgery needed if appropriate



#### Future Directions

- More long-term prospective pain and functional studies
- Photodynamic nails vs screw and cement constructs or both
- Utility of ablation on durability of constructs
- Similar constructs for long bones
- Timing of radiation
- Wider adoption outside of tertiary care facilities





#### References

- DiCaprio MR, Murtaza H, Palmer B, Evangelist M. Narrative review of the epidemiology, economic burden, and societal impact of metastatic bone disease. *Ann Joint*. 2022;7:28-28.
- English DI, Lea WB, King DM, Tutton SM, Neilson JC. Minimally Invasive Stabilization with or without Ablation for Metastatic Periacetabular Tumors. J Bone Joint Surg Am. 2021 Jul 7;103(13):1184-1192. doi: 10.2106/JBJS.20.00546. PMID: 34038393.
- Lee FY, Latich I, Toombs C, Mungur A, Conway D, Alder K, Ibe I, Lindskog D, Friedlaender G. Minimally Invasive Image-Guided Ablation, Osteoplasty, Reinforcement, and Internal Fixation (AORIF) for Osteolytic Lesions in the Pelvis and Periarticular Regions of Weight-Bearing Bones. J Vasc Interv Radiol. 2020 Apr;31(4):649-658.e1. doi: 10.1016/j.jvir.2019.11.029. Epub 2020 Mar 3. Erratum in: J Vasc Interv Radiol. 2020 Jun;31(6):1034. doi: 10.1016/j.jvir.2020.04.019. PMID: 32139256.
- Hirase T, Rowan C, Jacob T, Vemu SM, Aflatooni JO, Patel SS, Satcher RL, Lin PP, Moon BS, Lewis VO, Marco RAW, Bird JE. Percutaneous Cementation for Improvement of Pain and Function for Osteolytic Pelvic Metastasis: A Systematic Review. J Am Acad Orthop Surg. 2024 Jul 15;32(14):e726-e736. doi: 10.5435/JAAOS-D-23-00969. Epub 2024 Apr 30. PMID: 38709827.
- Lozano-Calderon SA, Clunk MJ, Gonzalez MR, Sodhi A, Krueger RK, Gruender AC, Greenberg DD. Assessing Pain and Functional Outcomes of Percutaneous Stabilization of Metastatic Pelvic Lesions via Photodynamic Nails: A Bi-Institutional Investigation of Orthopaedic Outcomes. JB JS Open Access. 2024 Jul 10;9(3):e23.00148. doi: 10.2106/JBJS.OA.23.00148. PMID: 38988331; PMCID: PMC11233101.
- Lea WB, Tutton SM, Alsaikhan N, et al. Pelvis weight-bearing ability after minimally invasive stabilizations for periacetabular lesion. *Journal Orthopaedic Research*. 2021;39(10):2124-2129.

CITY OF HOPE

32

### Questions?

■ Thank you to Dr. Jonathon Kessler and Dr. Alana Munger for collaboration